

Table 2. Number, median days, incidence rate¹ and relative standard errors of nonfatal occupational injuries and illnesses with days away from work² involving musculoskeletal disorders³ by selected parts of body, Oklahoma, 2002

Part of body		Number	Median days away from work	Incidence rate	Relative standard error
Total		5,961	11	60.5	6.0
1	Neck, Including Throat	141	19	1.4	19.8
10	Neck, except internal location of diseases or disorders	141	19	1.4	19.8
2	Trunk	4,088	8	41.5	6.3
21	Shoulder, including clavicle, scapula	622	21	6.3	10.5
22	Chest, including ribs, internal organs	--	--	--	--
220	Chest, except internal location of diseases or disorders	--	--	--	--
23	Back, including spine, spinal cord	2,546	6	25.8	6.9
230	Back, including spine, spinal cord, unspecified	1,106	8	11.2	8.6
231	Lumbar region	1,390	6	14.1	8.0
232	Thoracic region	--	--	--	--
238	Multiple back regions	17	3	0.2	54.6
24	Abdomen	716	14	7.3	9.9
240	Abdomen, except internal location of diseases or disorders	108	7	1.1	22.5
241	Internal abdominal location, unspecified	110	30	1.1	22.2
242	Stomach organ	--	--	--	--
245	Intestines, peritoneum	467	14	4.7	11.7
2450	Intestines, peritoneum, unspecified	465	14	4.7	11.7
2451	Peritoneum	--	--	--	--
25	Pelvic region	149	18	1.5	19.3
250	Pelvic region, unspecified	--	--	--	--
251	Hip(s)	56	38	0.6	30.7
254	Groin	91	14	0.9	24.4
28	Multiple trunk locations	--	--	--	--
3	Upper extremities	916	17	9.3	9.1
31	Arm(s)	305	4	3.1	14.0
310	Arm(s), unspecified	125	2	1.3	20.9
311	Upper arm(s)	--	--	--	--
312	Elbow(s)	94	9	1.0	23.9
313	Forearm(s)	19	7	0.2	52.1
318	Multiple arm(s) locations	44	17	0.5	34.6
32	Wrist(s)	438	36	4.4	12.0
33	Hand(s), except finger(s)	60	31	0.6	29.7
34	Finger(s), fingernail(s)	51	42	0.5	32.2
38	Multiple upper extremities locations	63	15	0.6	29.0
382	Hand(s) and wrist(s)	19	26	0.2	52.8
383	Hand(s) and arm(s)	--	--	--	--
389	Multiple upper extremities locations, n.e.c.	--	--	--	--
4	Lower extremities	524	40	5.3	11.2
41	Leg(s)	503	40	5.1	11.4
410	Leg(s), unspecified	--	--	--	--
411	Thigh(s)	40	3	0.4	36.1
412	Knee(s)	454	40	4.6	11.8
413	Lower leg(s)	--	--	--	--
419	Leg(s), n.e.c.	--	--	--	--
42	Ankle(s)	--	--	--	--
43	Foot(feet), except toe(s)	--	--	--	--
430	Foot(feet), except toe(s), unspecified	--	--	--	--
8	Multiple Body Parts	278	10	2.8	14.6
9	Other Body Parts	15	3	0.2	58.8
99	Other body parts, n.e.c.	15	3	0.2	58.8

Table 2. Number, median days, incidence rate¹ and relative standard errors of nonfatal occupational injuries and illnesses with days away from work² involving musculoskeletal disorders³ by selected parts of body, Oklahoma, 2002

Part of body		Number	Median days away from work	Incidence rate	Relative standard error
999	Other body parts, n.e.c.	15	3	0.2	58.8
9999	Nonclassifiable	15	3	0.2	58.8

¹ Incidence rates represent the number of injuries and illnesses per 10,000 full-time workers and were calculated as: $(N / EH) \times 20,000,000$ where,

N = number of injuries and illnesses,
EH = total hours worked by all employees during the calendar year,
20,000,000 = base for 10,000 full-time equivalent workers (working 40 hours per week, 50 weeks per year).

² Days away from work include those which result in days away from work with or without job transfer or restriction.

³ Includes cases where the nature of injury is: sprains, strains, tears; back pain, hurt back; soreness, pain, hurt, except back; carpal tunnel syndrome; hernia; or musculoskeletal system and connective tissue diseases and disorders and when the event or exposure leading to the injury or illness is: bodily reaction/bending, climbing, crawling, reaching, twisting; overexertion; or repetition. Cases of Raynaud's phenomenon, tarsal tunnel syndrome, and herniated spinal discs are not included. Although these cases may be considered MSD's, the survey classifies these cases in categories that also include non-MSD cases.

NOTE: Dashes indicate data that do not meet publication guidelines or data for incidence rates less than .05 per 10,000 full-time workers. The scientifically selected probability sample used was one of many possible samples, each of which could have produced different estimates. A measure of sampling variability for each estimate is available upon request.

SOURCE: Bureau of Labor Statistics, U.S. Department of Labor, April 2004